

List of publications related to IBEX collagen biomarker assays

1. Ameye LG, Deberg M, Oliveira M, Labasse A, Aeschlimann JM, Henrotin Y. (2007). The chemical biomarkers C2C, Coll2-1 and Coll2-1NO2 provide complimentary information on type II collagen catabolism in healthy and osteoarthritic mice. *Arthritis Rheum* 56: 3336-46.
2. Antoniou J, Steffen T, Nelson F, Winterbottom N, Hollander AP, et al. (1996). The human lumbar intervertebral disc. Evidence for changes in the biosynthesis and denaturation of the extracellular matrix with growth, maturation, ageing and degeneration. *J Clin Invest* 98: 996-1003.
3. Armstrong L, Thickett DR, Mansell JP, Ionescu M, Hoyle E, et al. (1999). Changes in collagen turnover in early acute respiratory distress syndrome. *Am J Resp Critical Care Med* 160: 1910-15.
4. Aurich M, Squires GR, Reiner A, Mollenhauer JA, Keuttner KE et al (2005). Differential matrix degradation and turnover in early cartilage lesions of human knee and ankle joints. *Arthritis Rheum* 52: 112-19.
5. Bay-Jensen AC, Andersen TL, Charni-Ben Tabassi N, Kristensen PW et al. (2008). Biochemical markers of type II collagen breakdown and synthesis are positioned at specific sites in human osteoarthritic knee cartilage. *Osteoarthritis Cartilage* 16; 615-23.
6. Billingham RC, Dahlberg L, Ionescu M, Reiner A, Bourne R et al. (1997). Enhanced cleavage of type II collagen by collagenases in osteoarthritic articular cartilage. *J Clin Invest* 99:1534-45.
7. Billingham RC, Wu W, Ionescu M, Reiner A, Dahlberg L, Chen J, Van Wart H, Poole AR. (2000) Comparison of the degradation of type II collagen and proteoglycan in nasal and articular cartilage induced by interleukin-1 and the selective inhibition of type II collagen cleavage by collagenase. *Arthritis Rheum.* 43: 664-672.
8. Billingham RC, Brama PA, van Weeren PR, Knowlton MS, McIlwraith CW. (2004). Evaluation of serum concentrations of biomarkers of skeletal metabolism and results of radiography as indicators of severity of osteochondrosis in foals. *Am J Vet Res.* 65:143-50.
9. Boeth H, MacMahon A, Poole AR, Buttgerit F, Onnefjord P et al. (2017). Differences in biomarkers of cartilage matrix turnover and their changes over 2 years in adolescent and adult volleyball athletes. *J Exp Orthop* 4: 1-11.
10. Briot K, Roux C, Gossec L, Charni N, Kolta S et al. (2008). Effects of etanercept on serum biochemical marker of cartilage metabolism in patients with spondyloarthritis. *J Rheumatol* 35:310-14
11. Cahue S, Sharma L, Dunlop D, Ionescu M, Song J. et al. (2007). The ratio of type II collagen breakdown to synthesis and its relationship with the progression of knee osteoarthritis. *Osteoarthritis Cartilage* 15: 819-23.
12. Carey DE, Alini M, Ionescu M, Hyams JS, Rowe JC, et al. (1997). Serum content of the C-propeptide of the cartilage molecule type II collagen in children. *Clin Exp Rheumatol* 15: 325-8.
13. Celeste C, Ionescu M, Poole AR, Laverty S. (2005). Repeated intraarticular injections of triamcinolone acetonide alter cartilage matrix metabolism measured by biomarkers in synovial fluid. *J Orthop Res* 23:602-10.

14. Chandran V, Cook RJ, Edwin J, Shen H, Pellett FJ et al. (2010). Soluble biomarkers differentiate patients with psoriatic arthritis from those with psoriasis without arthritis. *Rheumatology* 49:1399-405.
15. Chu Q, Lopez M, Hayashi K, Ionescu M, Billingham RC et al. (2002). Elevation of a collagenase generated type II collagen neoepitope and a proteoglycan epitope in synovial fluid following induction of joint instability in the dog. *Osteoarthritis Cartilage* 10: 662-9.
16. Chu CR, Sheth S, Erhart-Hledik JC, Do B, Titchenal, MR, Andriacchi, TP. (2017). Mechanically stimulated biomarkers signal cartilage changes over 5 years consistent with disease progression in medial knee osteoarthritis patients. *J Orthop Res*. 2017 Sep 1. doi: 10.1002/jor.23720. [Epub ahead of print].
17. Cibere J, Thorne A, Kopec JA, Singer J, Canvin J, et al. (2005). Glucosamine sulfate and cartilage : type II collagen degradation in patients with knee osteoarthritis: randomized discontinuation trial results employing biomarkers. *J Rheumatol* 32: 896-902.
18. Cibere J, Zhang H, Garnero P, Poole AR, Lobanok T. et al. (2009). Association of biomarkers with with pre-radiographically defined and radiographically defined knee osteoarthritis in a population-based cohort. *Arthritis Rheum* 60: 1372-80.
19. Conrozier T, Poole AR, Ferrand F, Mathieu P, Vincent F et al. (2008). Serum concentrations of type II collagen biomarkers (C2C, C1,2C and CPII) suggest different pathophysiologicals in patients with hip osteoarthritis. *Clin. Exp. Rheumatol*. 26: 430-5.
20. Conrozier T, Ferrand F, Poole AR, Verret C, Mathieu P, Ionescu M, Vincent F, Piperno M, Spiegel A, Vignon E. (2007) Differences in biomarkers of type II collagen in atrophic and hypertrophic osteoarthritis of the hip: implications for the differing pathobiologies. *Osteoarthritis Cartilage* 15: 462-7.
21. Dahlberg L, Billingham C, Manner P, Ionescu M, Reiner A et al. (2000). Selective enhancement of collagenase-mediated cleavage of resident type II collagen in cultured osteoarthritic cartilage and arrest with a synthetic inhibitor that spares collagenase 1 (matrix metalloproteinase 1). *Arthritis Rheum* 43: 673-82.
22. Dejica VM, Mort JS, Laverty S, Percival MD, Antoniou J et al. (2008). Cleavage of type II collagen by cathepsin K in human osteoarthritis cartilage. *Am J Pathol* 173: 161-9.
23. Dejica V, Mort JS, Laverty S, Antoniou J, Zukor DJ et al. (2012). Increased type II collagen cleavage by cathepsin K and collagenase activities with aging and osteoarthritis in human articular cartilage. *Arthritis Res Therapy* 14: R113.
24. Donabedian M, van Weeren PR, Perona G, Fleurance G, Robert C et al. (2008). Early changes in biomarkers of skeletal metabolism and their association to the occurrence of osteochondrosis (OC) in the horse. *Equine Vet J* 40:253-9.
25. de Grauw JC, Donabedian M, van der Lest CH, Perona G, Robert C et al. (2011). Assessment of synovial fluid biomarkers in healthy foals and in foals with tarsocrural osteochondrosis. *Vet J* 190:390-5.
26. Engstrom-Laurent A. (1989). Changes in hyaluronan concentration in tissues and body fluids in disease states. In: *The Biology of Hyaluronan*. D. Evered, J. Whelan (Eds.), John Wiley and Sons, Chichester, U.K., pp. 233-247.
27. Fernandez-Moreno M, Soto-Hermida A, Oreiro N, Pertega S, Fenandez-Lopez C et al. (2012). Mitochondrial haplogroups define two phenotypes of osteoarthritis. *Front Physiol* 3:129
28. Fraser A, Fearon U, Billingham RC, Ionescu M, Reece R et al. (2003). Turnover of type II collagen and aggrecan in cartilage matrix at the outset of inflammatory arthritis in humans. *Arthritis Rheum*. 48: 3085-95.

29. Frisbie DD, Ray CS, Ionescu M, Poole AR, Chapman PL, McIlwraith CW. (1999). Measurement of synovial fluid and serum concentrations of the 846 epitope of chondroitin sulfate and of carboxy peptides of type II procollagen for diagnosis of osteochondral fragmentation in horses. *Am J Vet Res* 60:306-9.
30. Frisbie DD, Al-Sobayil F, Billingham RC, Kawcak CE, McIlwraith CW. (2008). Changes in synovial fluid and serum biomarkers with exercise and early osteoarthritis in horses. *Osteoarthritis Cartilage*. 16:1196-2004.
31. Glant TT, Mikecz PJ, Roughley PJ, Buzas E, Poole AR. (1986) Age related changes in protein-related epitopes of human articular cartilage proteoglycans. *Biochem J*. 236: 71-75.
32. Goode AP, Marshall SW, Kraus VB, Renner JB, Stürmer T et al. (2012). Associations between serum and urine biomarkers and lumbar spine individual radiographic features: the Johnston County osteoarthritis project. *Osteoarthritis Cartilage* 20: 1286-93.
33. Hatcher CC, Collins AT, Kim SY, Michel LC, Mostertz WC 3rd et al. (2017). Relationship between T1rho magnetic resonance imaging, synovial fluid biomarkers, and the biochemical and biomechanical properties of cartilage. *J Biomech* ;55:18-26.
34. Hayashi PA, Kim SY, Lansdowne JL, Kapatkin A, Dejardin LM. (2009). Evaluation of a collagenase generated osteoarthritis biomarker in naturally occurring canine cruciate disease. *Vet Surg* 38: 1117-21
35. He G, Chen X, Zhang G, Lin H, Wu X. (2014). Detection of urine C2C and trace element level in patients with knee osteoarthritis. *Cell Biochem Biophys* 70:475-9.
36. Hollander A P, Heathfield T F, Webber C, Iwata Y, Bourne R, Rorabeck C, Poole A R. (1994) Increased damage to type II collagen in osteoarthritic articular cartilage detected by a new immunoassay. *J Clin Invest* 93: 1722-32
37. Huebner JL, Kraus VB. (2006), Assessment of the utility of biomarkers of osteoarthritis in the guinea pig. *Osteoarthritis Cartilage*. 14: 923-30.
38. Hunt MA, Pollock C L, Kraus VB, Saxne T, Peters S et al. (2013). Relationships amongst osteoarthritis biomarkers, dynamic knee joint load, and exercise: results from a randomized controlled pilot study. *BMC Musculoskeletal Disord* 14:115.
39. Ishijima M, Watari T, Naito K, Kaneko H, Futami I et al. (2011). Relationships between biomarkers of cartilage, bone, synovial metabolism and knee pain provide insights into the origins of pain in early knee osteoarthritis. *Arthritis Res Therapy* 13: R22
40. Jansen NW, Roosendaal G, Lundin B, Heijnen L, Mause-Bunschoten E et al. (2009). The combination of the biomarkers urinary C-terminal telopeptide of type II collagen, serum cartilage oligomeric matrix protein, and serum chondroitin sulfate 846 reflects cartilage damage in hemophilic arthropathy. *Arthritis Rheum* 60: 290-8.
41. Kilgallon CP, Larsen S, Wong A, Yellowley C. (2015). Analysis of a collagen II degradation protein C2C and a collagen II formation protein CPII in serum of Asian elephants (*Elephas maximus*). *J Zoo Wild Led* 46: 146-9.
42. Kim T-H, Stone M, Payne U, Zhang X, Ionescu M et al. (2005). Cartilage biomarkers in ankylosing spondylitis. Relationship to clinical variables and treatment response. *Arthritis Rheum* 52: 885-91.
43. King KB, Lindsey CT, Dunn TC, Ries MD, Steinbach LS, Majumdar S. (2004). A study of the relationship between molecular biomarkers of joint degeneration and the magnetic resonance-measured characteristics of cartilage in 16 symptomatic knees. *Magn. Reson. Imaging* 22: 1117-23.

44. Kojima T, Mwale F, Yasuda T, Girard C, Poole AR, Lavery S. (2001). Early degradation of type IX and type II collagen with the onset of experimental arthritis. *Arthritis Rheum* 44: 120-7.
45. Kojima T, Kojima M, Noda K, Ishiguro N, Poole AR. (2008). Influences of menopause, aging and gender on the cleavage of type II collagen in cartilage in relationship to bone turnover. *Menopause* 15: 133-7.
46. Kong SY, Stabler TV, Criscione LG, Elliott AL, Jordan KM, Kraus VB. (2006). Diurnal variation of serum and urine biomarkers in patients with radiographic knee osteoarthritis. *Arthritis Rheum* 54: 2496-504
47. Kraus VB, Burnett B, Coindreau J, Cottrell S, Eyre D et al. (2011). Application of biomarkers in the development of drugs intended for the treatment of osteoarthritis. *Osteoarthritis Cartilage* 19: 515-42
48. Kraus VB, Collins JE, Hargrove D, Losina E, Nevitt M et al. (2016) Predictive validity of biochemical biomarkers in knee osteoarthritis: data from the FNIH OA Biomarkers Consortium. *Ann Rheum Dis*. 76:186-95.
49. Kubomura D, Ueno T, Yamada M, Tomonaga M, Nagaoka I. (2017). Effect of N-acetylglucosamine administration on cartilage metabolism and safety in healthy subjects without symptoms of arthritis: A case report. *Exp. Ther. Med.* 13: 1614-21.
50. Kumahashi N, Sward P, Larsson S, Lohmander LS, Frobell R, Struglics A. (2015). Type II collagen C2C epitope in human synovial fluid and serum after knee injury-associations with molecular and structural markers of injury. *Osteoarthritis Cartilage* 23: 1506-12.
51. Lavery S, Ionescu M, Marcoux M, Boure L, Doize B, Poole AR. (2000). Alterations in cartilage type II procollagen and aggrecan contents in synovial fluid in equine osteochondrosis. *J Orthop Res* 18: 399-405.
52. Lavery S, O’Kouneff S, Ionescu M, Reiner A, Pidoux I et al (2002). Excessive degradation of type II collagen in articular cartilage in equine osteochondrosis. *J Orthop Res* 20: 1282-9.
53. Leff RL, Elias I, Ionescu M, Reiner A, Poole AR. (2003). Molecular changes in human osteoarthritis cartilage after 3 weeks of oral administration of BAY12-9566, a matrix metalloproteinase inhibitor. *J Rheumatol* 30: 544-9.
54. Lettry V, Sumie Y, Mitsuda K, Tagami M, Hosoya K et al. (2010). Divergent diagnosis from arthroscopic findings and identification of CPII and C2C for detection of cartilage degradation in horses. *Jpn J Vet Res* 57: 197-206.
55. Lohmander SL, Yoshihara Y, Roos H, Kobayashi T, Yamada H, Shinmei M. (1996) Procollagen II C-Propeptide in Joint fluid: Changes in concentration with age, time after knee injury and osteoarthritis. *J. Rheumatol.* 23: 1765-1769.
56. Lohmander LS, Ionescu M, Jugessur H, Poole AR. (1999). Changes in joint cartilage aggrecan after knee injury and in osteoarthritis. *Arthritis Rheum.* 42: 534-544.
57. Ma J, Sorsa T, Billingham R, Poole AR, Kittu U et al. (2003) Direct evidence of collagenolysis in chronic periodontitis. *J Periodontal Res.* 38: 564-7.
58. Maksymowych WP, Poole AR, Hiebert L, Webb A, Ionescu M et al. (2005). Etanercept exerts beneficial effects on articular cartilage biomarkers of degradation and turnover in patients with ankylosing spondylitis. *J Rheumatol* 32: 1911-7.
59. Manicourt D-H, Azria M, Mindeholm L, Thonar E J-M, Devogelaer J-P. (2006). Oral salmon calcitonin reduces Lequesne’s algofunctional index scores and decreases urinary and serum levels of biomarkers of joint metabolism in knee osteoarthritis. *Arthritis Rheum.* 54: 3205-11.

60. Mansson B, Carey D, Alini M, Ionescu M, Rosenberg LC, Poole AR, Heinegard D, Saxne T. (1995). Cartilage and bone metabolism in rheumatoid arthritis. *J. Clin. Invest.* 95: 1071-1077.
61. Matyas JR, Atley L, Ionescu M, Eyre DR, Poole AR. (2004). Analysis of cartilage biomarkers in the early phase of canine experimental osteoarthritis. *Arthritis Rheum* 50: 543-52.
62. Mazzuca SA, Poole AR, Brandt KD, Katz BP, Lane KA, Lobanok T. (2006) Associations between joint space narrowing and molecular markers of collagen and proteoglycan turnover in patients with knee osteoarthritis. *J Rheumatol* 33: 1147-51.
63. Mitchell PG, Magna HA, Reeves LM, Lopresti-Morrow LL, Yocum SA et al. (1996). Cloning, expression and type II collagenolytic activity of matrix metalloproteinase-13 from human osteoarthritic cartilage. *J Clin Invest* 97: 761-8.
64. Mullan RH, Matthews C, Bresnihan B, Fitzgerald O, King L et al. (2007). Early changes in serum type II collagen biomarkers predict radiographic progression at one year in inflammatory arthritis patients after biologic therapy. *Arthritis Rheum* 56: 2919-28.
65. Mündermann A, Klenk C, Billich C, Nüesch C, Pagenstert G et al. (2017). Changes in cartilage biomarker levels during a transcontinental multistage footrace over 4486 km. *Am J Sports Med.* 2017 Jun1:363546517712945.doi:10.1177/0363546517712945. [Epub ahead of print]
66. Mwale F, Billingham C, Wu W, Alini M, Webber C, Reiner A, Ionescu M, Poole J, Poole A R. (2000) Selective assembly and remodelling of collagens II and IX associated with expression of the chondrocyte hypertrophic phenotype . *Developmental dynamics* 218(4):648-62.
67. Mwale F, Tchetina E, Wu W, Poole AR. (2002). The assembly and remodeling of the extracellular matrix in the growth plate in relationship to mineral deposition and cellular hypertrophy: an in situ study of collagens II and IX and proteoglycan. *J Bone Min Res* 17: 275-83.
68. Nelson F, Dahlberg L, Laverty S, Reiner A, Pidoux I et al. (1998). Evidence for altered synthesis of type II collagen in patients with osteoarthritis. *J Clin Invest* 102: 2115-25.
69. Nelson F, Billingham RC, Pidoux I, Reiner A, Langworthy M, et al. (2006). Early post-traumatic osteoarthritis-like changes in human articular cartilage following rupture of the anterior cruciate ligament. *Osteoarthritis Cartilage* 14:114-9.
70. Nemirovskiy OV, Dufield DR, Sunyer T, Aggarwal P, Welsch DJ, Mathews WR. (2007). Discovery and development of a type II collagen neoepitope (TIINE) biomarker for matrix metalloproteinase activity: from in vitro to in vivo. *Annal Biochem* 361: 93-101.
71. Niki Y, Takeuchi T, Nakayama M, Nagasawa H, Kurasawa T et al. (2012). Clinical significance of cartilage biomarkers for monitoring structural joint damage in rheumatoid arthritis patients treated with anti-TNF therapy. *PLoS One* 7: e37447
72. Oldenburg J, Zimmermann R, Katsarou O, Zanon E, Kellermann E et al. (2016). Potential biomarkers of haemophilic arthropathy: correlations with compatible additive magnetic resonance imaging scores. *Haemophilia* 22: 760-4.
73. Otterness IG, Swindell AC, Zimmerer RO, Poole AR, Ionescu M, Weiner E. (2000). An analysis of 14 molecular markers for monitoring osteoarthritis: segregation of the 14 markers into clusters and distinguishing osteoarthritis at baseline. *Osteoarthritis Cartilage* 8: 180-5.
74. Owens BD, Cameron KL, Bokshan SL, Clifton KB, Svoboda SJ, Wolf, JM. (2017). Serum cartilage biomarkers and shoulder instability. *Orthopedics* 40: 34-6.

75. Pietrosimone B, Blackburn JT, Harkey MS, Luc BA, Hackney AC et al. (2016a). Greater mechanical loading during walking is associated with less collagen turnover in individuals with anterior cruciate ligament reconstruction. *Am J Sports Med.* 44:425-32.
76. Pietrosimone B, Blackburn T, Harkey MS, Luc BA, Hackney AC et al. (2016b). Walking speed as a potential indicator of cartilage breakdown following anterior cruciate ligament reconstruction. *Arthritis Care Res (Hoboken).* 68: 793-800.
77. Pietrosimone B, Loeser RF, Blackburn JT, Padua DA, Harkey MS, et al. (2017). Biochemical markers of cartilage metabolism are associated with walking biomechanics six-months following anterior cruciate ligament reconstruction. *J Orthop Res* 2 MAR 2017, DOI: 10.1002/jor.23534
78. Poole AR, Ionescu M, Swan A, Dieppe PA. (1994). Changes in cartilage metabolism in arthritis are reflected by altered serum and synovial fluid levels of the cartilage proteoglycan aggrecan. Implications for pathogenesis. *J Clin Invest* 94: 25-33.
79. Poole AR, Alini M, Hollander AH. (1995) Cellular biology of cartilage degradation. In: Henderson B, Edwards, J.C.W, Pettipher, E.R, (Eds.). *Mechanisms and models in rheumatoid arthritis.* London: Academic Press; pp. 163-204.
80. Poole AR., Ionescu M, Fitzcharles MA, Billingham R. (2004). The assessment of cartilage degradation *in vivo*: Development of an immunoassay for the measurement in body fluids of type II collagen cleaved by collagenases. *J Immunological Methods*, 294:145-53.
81. Poole AR (2016). Current opinion: where are we in our understanding and treatment of osteoarthritis ? *Swiss Med Wkly* 146: w14340.
82. Poole AR, Ha N, Bourdon S, Sayre EC, Guermazi A, Cibere J. (2016). Ability of a urine assay of type II collagen cleavage by collagenases to detect early onset and progression of articular cartilage degeneration: Results from a population-based cohort study. *J Rheumatol.* 43:1864-70.
83. Prink A, Hayashi K, Kim SY, Kim J, Kapatkin A. (2010). Evaluation of a collagenase generated osteoarthritis biomarker in the synovial fluid from elbow joints of dogs with medial coronoid disease and unaffected dogs. *Vet Surg* 39: 65-70.
84. Ramonda R, Lorenzin M, Modesti V, Campana C, Ortolan A, et al. (2013). Serological markers of erosive hand osteoarthritis. *Eur J Intern Med* 24:11-5
85. Richette P, Eymard RP, Deberg M, Vidaud D, de Kerguenec C et al. (2010). Increase in type II collagen turnover after iron depletion in patients with hereditary haemochromatosis. *Rheumatology (Oxford)* 49: 760
86. Rizkalla G, Reiner A, Bogoch E, Poole AR. (1992). Studies of the articular cartilage proteoglycan in health and osteoarthritis: Evidence for molecular heterogeneity and extensive molecular changes in disease. *J Clin Invest* 90: 2268-77.
87. Robion FC, Doize B, Boure L, Marcoux M, Ionescu M et al. (2001). Use of synovial fluid markers of cartilage synthesis and turnover to study effects of repeated intra-articular administration of methyl prednisolone acetate on articular cartilage. *J Orthop Res* 19: 250-258.
88. Sharma L, Dunlop D, Ionescu M, Song J, Lobanok T, King L, Cahue S, Poole AR.(2007) The Ratio of Collagen Breakdown to Collagen Synthesis and its Relationship with the Progression of Knee Osteoarthritis. *Osteoarthritis Cartilage.* 15(7): 819-823.
89. Song X-Y, Zeng L, Jin W, Thompson J, Mizel DE et al. (1999). Secretory leucocyte protease inhibitor suppresses the inflammation and joint damage of bacterial cell wall-induced arthritis. *J Exp Med* 190: 535-42.

90. Soto-Hermida A, Fernandez-Moreno M, Pertaga-Diaz S, Oreiro N, Fernando-Lopez C et al. (2015) Mitochondrial DNA haplogroups modulate the radiographic progression of Spanish patients with osteoarthritis. *Rheumatol. Int* 35: 337-44.
91. Squires G, Okouneff S, Ionescu M, Poole AR.(2003). Pathobiology of focal lesion development in aging human articular cartilage reveals molecular matrix changes characteristic of osteoarthritis. *Arthritis Rheum* 48: 1261-70.
92. Stoop R, van der Kraan PM, Buma P, Hollander AP, Billingham RC, et al. (1999). Type II collagen degradation in spontaneous osteoarthritis in C57B1/6 and BALB/c mice. *Arthritis Rheum* 42: 2381- 9.
93. Stoop R, Buma P, van der Kraan P, Hollander AP, Billingham RC et al. (2001). Type II collagen degradation in articular fibrillation and anterior cruciate ligament transection in rats. *Osteoarthritis Cartilage* 9: 308-15.
94. Sugiyama S, Itokazu M, Suzuki Y, Shimizu K. (2003) Procollagen II C propeptide level in the synovial fluid as a predictor of radiographic progression in early knee osteoarthritis. *Ann Rheum Dis* 62: 27-32.
95. Sukhova GK, Schönbeck Y, Rabkin E, Schoen FJ, Poole AR, Billingham RC, Libby P. (1999) Evidence for increased collagenolysis by interstitial collagenases -1 and -3 in vulnerable human atheromatous plaques. *Circulation* 99:2503-2509.
96. Svoboda SJ, Harvey TM, Owens BD, Brechue WF, Tarwater PM, Cameron KL. (2013). Changes in serum biomarkers of cartilage turnover after anterior cruciate ligament injury. *Am J Sports Med* 41: 2108-16.
97. Svoboda, SJ, Owens, BD, Harvey TM, Tarwater,PM, Brechue WF Cameron, KL. (2016). Biomarkers of collagen turnover and subsequent anterior cruciate ligament rupture. *Am J Sports Med.* 44: 1687-93
98. Tamm AO, Kumm J, Tamm A, Lintrop M, Kukner A et al. (2014). Cartilage collagen neoepitope C2C and clinical parameters in middle-aged patients with knee problems. Correlations of urinary output of C2C with cartilage lesions, KOOS values and functional abilities of lower limb. *Osteoarthritis Cartilage* 22: S70-S71.
99. Tomonaga A, Takahashi T, Tanaka YT, Tsuboi M, Ito K, Nagaoka I. (2017). Evaluation of the effect of salmon nasal proteoglycan on biomarkers for cartilage metabolism in individuals with knee joint discomfort: A randomized double-blind placebo-controlled clinical study. *Exp Ther Med.* 14:115-26.
100. Tourville TW, Johnson RJ, Slauterbeck JR, Naud S, Beynon BD. (2013). Relationship between markers of type II collagen metabolism and tibiofemoral joint space width changes after ACL injury and reconstruction. *Am J Sports Med* 41:779-87.
101. Trumble TN, Billingham RC, McIlwraith CW. (2003) Increased levels of synthesis and degradation markers of type II collagen in the synovial fluid of canine cruciate transection model of osteoarthritis . *Trans ORS* 27:0679.
102. Trumble TN, Scarborough AB, Brown MP. (2009). Osteochondral injury increases type II collagen degradation products (C2C) in synovial fluid of Thoroughbred racehorses. *Osteoarthritis Cartilage* 17:371-4.
103. Tyler JA. (1985).Articular cartilage cultured with catabolin (pig interleukin 1) synthesizes a decreased number of normal proteoglycan molecules. *Biochem J* 227: 869-78.
104. van der Rest M, Rosenberg LC, Olsen BR, Poole AR. (1986). Chondrocalcin is identical with the C-propeptide of type II collagen. *Biochem J* 237: 923-5.
105. van Meurs J, van Lent P, Stoop R, Holthuysen A, Singer I et al.(1999). Cleavage of aggrecan at the ASN³⁴¹-PHE³⁴² site coincides with the initiation of collagen damage in

- murine antigen-induced arthritis. A pivotal role for stromelysin 1 in matrix metalloproteinase activity. *Arthritis Rheum* 42: 2074-84.
106. van Spil WE, Jansen NW, Bijlsma JW, Reijman M, Degroot J et al. (2012). Clusters within a wide spectrum of biochemical markers of osteoarthritis: data from CHECK, a large cohort of individuals with very early symptomatic osteoarthritis. *Osteoarthritis Cartilage* 20: 745-54.
 107. van Vulpen LF, van Meergeren ME, Roosendaal G, Jansen NW, van Laar JM et al. (2015). Biochemical markers of joint damage increase shortly after a joint bleed; an explorative human and canine in vivo study. *Osteoarthritis Cart* 23: 63-9.
 108. Verstappen SMM, Poole AR, Ionescu M, King LE, Abrahamowicz M et al. (2006). Radiographic joint damage in rheumatoid arthritis is associated with differences in cartilage turnover and can be predicted by serum biomarkers: an evaluation from 1 to 4 years after diagnosis. *Arthritis Res Therapy* 8: R31.
 109. Vilar JM, Rubio M, Spinella G, Cuervo B, Sopena J, Cugat R et al. (2016) Serum collagen Type II cleavage epitope and serum hyaluronic acid as biomarkers for treatment monitoring of dogs with hip osteoarthritis. *PloS ONE* 11(2): e0149472. doi:10.1371/journal.pone.0149472.
 110. Wasilko SM, Tourville, TW, DeSarno MJ, Slauterbeck JR, Johnson RJ et al. (2016). Relationship between synovial fluid biomarkers of articular cartilage metabolism and the patient's perspective of outcome depends on the severity of articular cartilage damage following ACL trauma. *J Orthop Res* 34:820-7.
 111. Wei L, Fleming BC, Sun X, Teeple E, Wu W et al. (2010). Comparison of differential biomarkers of osteoarthritis with and without posttraumatic injury in the Hartley guinea pig model. *J Orthop Res* 28:900-6.
 112. Wu W, Billingham RC, Pidoux I, Antoniou J, Zukor D et al. (2002). Sites of collagenase cleavage and denaturation of type II collagen in aging and osteoarthritic articular cartilage and their relationship to the distribution of matrix metalloproteinase 1 and matrix metalloproteinase 13. *Arthritis Rheum* 46: 2087-94.
 113. Yamaguchi S, Aoyama T, Ito A, Nagai M, Iijima H et al. (2013). Effects of exercise level on biomarkers in a rat knee model of osteoarthritis. *J Orthop Res* 31:1026-31.
 114. Yoshida H, Kojima T, Kurokouchi K, Hanamura H et al. (2013). Relationship between pre-radiographic cartilage damage following anterior cruciate ligament injury and biomarkers of cartilage turnover in clinical practice: a cross-sectional observational study. *Osteoarthritis Cartilage* 21:831-8.
 115. Zack DJ, Bolon B, Hu Y-L, Poole AR, Feige U. (2003) Destruction of joint cartilage in rats with adjuvant-induced arthritis is reduced in an additive fashion by combination therapy with inhibitors of IL-1 and TNF. *Arthritis and Rheumatism* 48(S):S556